

Eisai Inc./Pfizer Inc.	IFC, 1
Forest Laboratories Inc. Namenda	16a-b
Janssen Pharmaceutica Eldercare	33
Razadyne	9-10
Risperdal	48, IBC, OBC
Medicine-On-Time	3
Merck Co., & Inc.	18
NADONA/LTC	27
PALs Software	48
QS/1, PrimeCare®	48
Wyeth Pharmaceuticals	5



PALs Software

- Designed for Assisted Living
- Comprehensive
- Affordable

“It’s brilliant...”

“It does everything!”

“PALs gives me more time with my residents.”

Includes Marketing, MARs, Assessments & Care Plans, QI, Resident Agreements & Service documentation, Billing, P&P. Tracks Supervision Dates, TO’s, Staff Training, plus much more.

Modules priced from \$500

Download FREE trial at:
www.PALsSoftware.com

Or Call for Online Demo
Toll Free: 1-877-724-7257

<h2 style="margin: 0;">RISPERDAL®</h2> <p style="margin: 0;">(RISPERIDONE) TABLETS/ORAL SOLUTION</p>	<h2 style="margin: 0;">RISPERDAL® M-TAB®</h2> <p style="margin: 0;">(RISPERIDONE) ORALLY DISINTEGRATING TABLETS</p>
--	---

BEFORE PRESCRIBING, PLEASE CONSULT COMPLETE PRESCRIBING INFORMATION OF WHICH THE FOLLOWING IS A BRIEF SUMMARY.

Increased Mortality in Elderly Patients with Dementia-Related Psychosis
Elderly patients with dementia-related psychosis treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo. Analyses of seventeen placebo controlled trials (modal duration of 10 weeks) in these patients revealed a risk of death in the drug-treated patients of between 1.6 to 1.7 times that seen in placebo-treated patients. Over the course of a typical 10 week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. RISPERDAL® (risperidone) is not approved for the treatment of patients with Dementia-Related Psychosis.

INDICATIONS AND USAGE
RISPERDAL® (risperidone) is indicated for the treatment of schizophrenia.
Monotherapy: RISPERDAL® is indicated for the short-term treatment of acute manic or mixed episodes associated with Bipolar I Disorder.
Combination Therapy: The combination of RISPERDAL® with lithium or valproate is indicated for the short-term treatment of acute manic or mixed episodes associated with Bipolar I Disorder.

CONTRAINDICATIONS
RISPERDAL® (risperidone) is contraindicated in patients with a known hypersensitivity to the product.

WARNINGS
Increased Mortality in Elderly Patients with Dementia-Related Psychosis
Elderly patients with dementia-related psychosis treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo. RISPERDAL® (risperidone) is not approved for the treatment of dementia-related psychosis (see Boxed Warning).

Neuroleptic Malignant Syndrome (NMS) A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with antipsychotic drugs. If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reinstitution of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

Tardive Dyskinesia A syndrome of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with antipsychotic drugs. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown. If signs and symptoms of tardive dyskinesia appear in a patient on RISPERDAL®, drug discontinuation should be considered. However, some patients may require treatment with RISPERDAL® despite the presence of the syndrome.

Cerebrovascular Adverse Events, Including Stroke, in Elderly Patients With Dementia-Related Psychosis
Cerebrovascular adverse events (e.g., stroke, transient ischemic attack), including fatalities, were reported in patients (mean age 85 years; range 73-97) in trials of risperidone in elderly patients with dementia-related psychosis. In placebo-controlled trials, there was a significantly higher incidence of cerebrovascular adverse events in patients treated with risperidone compared to patients treated with placebo. RISPERDAL® has not been shown to be safe or effective in the treatment of patients with dementia-related psychosis. (See also Boxed Warning, WARNINGS: Increased Mortality in Elderly Patients with Dementia-Related Psychosis.)

Hyperglycemia and Diabetes Mellitus Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including RISPERDAL®. Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment.

PRECAUTIONS
General

Orthostatic Hypotension: RISPERDAL® (risperidone) may induce orthostatic hypotension associated with dizziness, tachycardia, and in some patients, syncope, especially during the initial dose-titration period, probably reflecting its alpha-adrenergic antagonistic properties. Syncope was reported in 0.2% (6/2607) of RISPERDAL® treated patients in phase 2-3 studies. The risk of orthostatic hypotension and syncope may be minimized by limiting the initial dose to 2 mg total (either QD or 1 mg BID) in normal adults and 0.5 mg BID in the elderly and patients with renal or hepatic impairment (See DOSAGE AND ADMINISTRATION). Monitoring of orthostatic vital signs should be considered in patients for whom this is of concern. A dose reduction should be considered if hypotension occurs. RISPERDAL® should be used with particular caution in patients with known cardiovascular disease (history of myocardial infarction or ischemia, heart failure, or conduction abnormalities), cerebrovascular disease, and conditions which would predispose patients to hypotension e.g., dehydration and hypovolemia. Clinically significant hypotension has been observed with concomitant use of RISPERDAL® and antihypertensive medication.

Seizures: RISPERDAL® should be used cautiously in patients with a history of seizures.
Dysphagia: Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's dementia. RISPERDAL® and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia. (See also Boxed Warning, WARNINGS: Increased Mortality in Elderly Patients with Dementia-Related Psychosis.)

Osteodystrophy and Tumors in Animals: RISPERDAL® CONSTA™ produced osteodystrophy in male and female rats in a 1-year toxicity study and a 2-year carcinogenicity study at a dose of 40 mg/kg administered IM every 2 weeks. RISPERDAL® CONSTA™ produced renal tubular tumors (adenoma, adenocarcinoma) and adrenomedullary pheochromocytomas in male rats in the 2-year carcinogenicity study at 40 mg/kg administered IM every 2 weeks. In addition, RISPERDAL® CONSTA™ produced an increase in a marker of cellular proliferation in renal tissue in males in the 1-year toxicity study and in renal tumor-bearing males in the 2-year carcinogenicity study at 40 mg/kg administered IM every 2 weeks.

Hyperprolactinemia: As with other drugs that antagonize dopamine D₂ receptors, risperidone elevates prolactin levels and the elevation persists during chronic administration. Neither clinical studies nor epidemiologic studies conducted to date have shown an association between chronic administration of this class of drugs and tumorigenesis in humans; the available evidence is considered too limited to be conclusive at this time.

Potential for Cognitive and Motor Impairment: Somnolence was a commonly reported adverse event associated with RISPERDAL® treatment, especially when ascertained by direct questioning of patients. This adverse event is dose related. Patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that RISPERDAL® therapy does not affect them adversely.

Priapism: Rare cases of priapism have been reported.

Thrombotic Thrombocytopenic Purpura (TTP): A single case of TTP was reported in a 28-year-old female patient receiving RISPERDAL® in a large, open premarketing experience (approximately 1300 patients). She experienced jaundice, fever, and bruising, but eventually recovered after receiving plasmapheresis. The relationship to RISPERDAL® therapy is unknown.

Antiemetic Effect: Risperidone has an antiemetic effect in animals; this effect may also occur in humans, and may mask signs and symptoms of overdose with certain drugs or of conditions such as intestinal obstruction, Reye's syndrome, and brain tumor.

Body Temperature Regulation: Disruption of body temperature regulation has been attributed to antipsychotic agents. Caution is advised when prescribing for patients who will be exposed to temperature extremes.

Suicide: The possibility of a suicide attempt is inherent in schizophrenia, and close supervision of high risk patients should accompany drug therapy.

Use in Patients With Concomitant Illness: Clinical experience with RISPERDAL® in patients with certain concomitant systemic illnesses is limited. Patients with Parkinson's Disease or Dementia with Lewy Bodies who receive antipsychotics may be at increased risk of Neuroleptic Malignant Syndrome as well as having an increased sensitivity to antipsychotic medications. Manifestation of this increased sensitivity can include confusion, obtundation, postural instability with frequent falls, in addition to extrapyramidal symptoms. Caution is advisable in using RISPERDAL® in patients with diseases or conditions that could affect metabolism or hemodynamic responses.